

UNIVERSITÄTSKLINIK FÜR NEUROLOGIE

VORSTAND: O. UNIV. PROF. DR. WERNER POEWE  
BETREUER: A. UNIV. PROF. DR. GREGOR KARL WENNING

**DIE ROLLE DER KONVENTIONELLEN  
MAGNETRESONANZTOMOGRAPHIE  
IN DER DIFFERENTIALDIAGNOSE VON  
MULTISYSTEMATROPHIE UND  
MORBUS PARKINSON**

The Role of routine magnetic resonance imaging in the differential diagnosis of Multiple system atrophy  
and Parkinson's Disease

**Dissertation**

Zur Erlangung des Grades

**Doctor medicinae universae**

an der

Medizinischen Fakultät

der Leopold-Franzens-Universität Innsbruck

vorgelegt von Hannes Joachim Meier

Innsbruck, Januar 2002



## **Table of contents**

<b>Abstract</b> .....	3
<b>Zusammenfassung</b> .....	4
<b><u>1. Introduction</u></b> .....	5
1.1. Parkinson's disease (PD).....	6
1.1.1. Epidemiology.....	7
1.1.2. Aetiology and Pathogenesis.....	7
1.1.3. Clinical features and diagnosis.....	7
1.1.4. Therapy.....	8
1.2. Multiple system atrophy (MSA) .....	10
1.2.1. Epidemiology.....	10
1.2.2. Symptoms.....	11
1.2.3. Therapy.....	12
1.2.4. Diagnostic criteria for MSA.....	12
1.2.5. Additional investigations.....	16
1.3. MRI in PD and MSA.....	17.
<b><u>2. Patients and Methods</u></b> .....	19
2.1. Patients.....	19
2.2. MRI Evaluation.....	20
2.3. Statistical analysis .....	22

<b>3. Results</b> .....	23
3.1. Patients.....	23
3.2. Basal ganglia findings.....	23
3.3. Infratentorial abnormalities.....	25
3.4. Overall sensitivity, specificity and positive predictive value.....	26
3.5. Cortical atrophy.....	27
3.6. Clinical correlation.....	30
3.7. Proposed MRI criteria for MSA.....	31
<b>4. Discussion</b> .....	36
<b>5. References</b> .....	39